

ISOLATION AND SCREENING OF POTENTIAL BACTERIA WITH BIOFILM FORMATION ABILITY AND THEIR HYDROLYTIC ENZYMES FOR ENHANCED BIOGAS PRODUCTION

Nurul Alia Fazil, Md Zahangir Alam, Azlin Suhaida Azmi, Mariatul Fadzillah Mansor

Bioenvironmental Engineering Research Centre (BERC), Department of Biotechnology Engineering, Faculty of Engineering, International Islamic University Malaysia, P. O. Box 10, 50728 Kuala Lumpur, Malaysia
Email: aliafazil@live.iium.edu.my, zahangir@iium.edu.my, azlinsu76@iium.edu.my, mariatul@iium.edu.my

ABSTRACT

Biofilm mediated anaerobic digestion of POME are to be studied to produce biogas. Out of 120 strains isolated from palm oil mill effluent (POME), palm kernel cake (PKC), and food waste compost, only 13 strains showed biofilm producing ability with significant amount at $OD_{595nm} (>0.01)$. In hydrolytic enzyme assay test, which are amylolytic enzyme assay, cellulolytic enzyme assay, lipolytic enzyme assay, and proteolytic enzyme assay all strains does not secrete protease enzyme, and strain number 1C and 23 does not secrete any hydrolytic enzymes. Strain number 23C and 30C shows positive result for cellulase, amylase and lipase enzymes to be tested as single strain bacteria for prospect research on producing biogas while other strains will be treated as mixed culture in future research.

Keywords: Biofilm, hydrolytic enzyme, isolation, palm oil mill effluent.

INTRODUCTION

Malaysia is one of the biggest palm oil producer in the world. Parallel to palm oil production, high organic waste, palm oil mill effluent (POME) is also produced. Due to the high organic content, POME is to be treated prior to discharge to any water body. On the other hand, due to its high organic content, treatment of POME can be used to produce biogas, a type of renewable energy [1]. This opens the opportunity of turning waste to wealth.

Biogas production using POME as a feedstock has been studied by many researchers. Biogas production can be increased with better pre-treatment (digestion) [2 - 4]. Pre-treatment of feedstock includes mechanical, thermal, chemical and biological method. Commonly biological pre-treatment is much cheaper than others due to its energy saving (can operate at lower temperature, no aeration, etc.) but consumes much longer time. Biological pre-treatments comes in 3 mode; aerobic digestion, anaerobic digestion and enzyme addition. It is well known that enzyme is very effective but also is very costly. Aerobic digestion could lead to uncontrollable greenhouse gases emission [5] thus anaerobic digestion system was studied in order to efficiently collect biogas. Anaerobic granular sludge and biofilm system can overcome the problem of slow growth and process instability that occurs during anaerobic digestion [6].

Biofilm based system may be able to solve the problem of biomass washout because of the continuous mixing of effluent during anaerobic treatment of POME [7]. However, biofilm mediated anaerobic digestion of POME using single or multiple identified colonies of locally isolated bacteria has yet to be explored deeply. A study done by previous researcher have resulted in 28% increase in biogas production using biofilm mediated digestion from mixed culture of anaerobic sludge digester with liquid manure as the feedstock [8]. This study aims to isolate biofilm producing bacteria from POME and palm kernel cake (PKC) as the source of bacteria that is most related to the nature of POME and food waste compost as additional source of bacteria. The isolated bacteria then were studied for their ability to produce biofilm and hydrolytic enzyme secretion in order to understand their role in pre-treating POME for future research.

METHODS

Sample collection

Palm oil mill effluent (POME) and palm kernel cake (PKC) sample were collected at Sime Darby Sdn. Bhd. Banting, Selangor. Food waste compost was collected from composter in International Islamic University Malaysia (IIUM). Then the collected samples were taken for bacterial isolation.

Bacteria Isolation

One ml of POME sample was cultivated overnight in 10 ml of Luria Bertani (LB) broth at 37°C, and 150 rpm agitation. PKC and food compost waste were cultivated overnight by adding 0.1 gram of the sample into 100 ml LB broth and incubated at 37°C with 150 rpm agitation. The culture was diluted in LB broth up to 10^{-6} dilution factor and plated on LB agar (10^{-3} to 10^{-6}) and then incubated at 37°C for 24 to 48 hour. Any visible colony was transferred to the new fresh plate. Isolates from POME were labelled with numbers, isolates from PKC were labelled with numbers and letter P and isolates from food waste compost were labelled with numbers and letter C.

Biofilm Producer Screening

The isolated bacteria undergoes biofilm assay screening. Method of biofilm assay test was adapted from Djordjevic (2002) [9]. 5 ml of overnight cultures were prepared for each isolates and incubated at 37°C with 150 rpm shaking. The overnight cultures were diluted by adding 0.1 ml of culture into 10 ml LB broth and then mixed. The diluted cultures then were liquated 100 µl into each wells of sterile 96 wells microtiter plate, triplicated for each culture. After 24 hour and 48 hour of incubation, the growth of bacteria were taken at optical density (OD) 595 nm with 96 wells microtiter plate reader.

The plate then washed 5 times with sterile distilled water, and dried for 45 minutes. Each well of the plate then stained with 150 µl of 1% crystal violet solution for 45 minutes then washed again with sterile distilled water 5 times. After drying for 45 minutes, the wells of the plate were destained with 200 µl 95% ethanol solution and read OD_{595 nm}. Positive biofilm producing bacteria were used for hydrolysis enzyme assay test. Four hydrolysis enzyme assay were cellulolytic enzyme assay, lipolytic enzyme assay, amylolytic enzyme assay, and proteolytic enzyme assay.

Hydrolytic Enzyme Screening

Carboxymethylcellulose (CMC) agar was prepared to detect cellulose enzyme. The formula of the CMC agar was adapted from Gohel et. al (2014) [10]. 1% of CMC and 2% of bacteriological agar were prepared. The isolated bacteria was streaked on CMC agar plate and incubated for 24 hour. After 24 hour, the plates were stained with 2 ml Gram's iodine solution for 5 minute, and then washed with distilled water.

Lipase enzyme was detected using phenol red agar with olive oil as the indicator (0.01% w/v phenol red, 0.1% v/v olive oil, 0.1% w/v calcium chloride, and 2% w/v agar). The bacteria were streaked on agar plate and incubated at 37°C for 24 hour and 48 hour [11]. Colour changes were observed.

Amylase was detected using starch medium. The formulation of the agar is as follow; 2 grams soluble starch, 1 gram peptone, 1 gram yeast extract, 20 grams agar, and 1 litre distilled water [12]. After 24 hour to 48 hour incubation, the plates were stained with 2 ml Gram's iodine solution and for 5 minutes and then washed with distilled water.

The method of testing the presence of protease enzyme was adopted from Vijayaraghavan et. al (2013) [13]. For 1 litre of distilled water, 5 grams of peptic digest of animal tissue, 1.5 grams of beef extract, 1.5 grams of yeast extract, 5 grams of calcium chloride, 15 grams of agar and 0.0015% w/v of bromocresol green (BCG) reagent were added. BCG dye was prepared adding 0.56% w/v succinic acid, 0.1% w/v sodium hydroxide, 0.028% w/v BCG dye and 1% Brij-35. The pH then adjusted to pH 4.15 and the solution must be stored at 2-8°C. Protease producing bacteria will be screened on by streaking on the prepared plate and incubated at 37°C for 48 hour. Any colour changes will be recorded.


RESULTS

Isolation and screening for biofilm formation


120 isolates were acquired from POME, PKC and food waste compost. Out of all isolates obtained, only 13 strains were chosen to undergo hydrolysis enzyme screening. Table 1 shows the isolates and the source of bacteria. The strains chosen have significant amount of biofilm spectrophotometry reading at optical density (OD) 595 nm (<0.01). Figure 1 (a) and (b) shows the results of bacterial growth and biofilm amount at 24 hour and 48 hour.

Table 1. Isolates and its sources.

Source	Isolates
POME	7, 11, 23, 36, 39
PKC	3P, 19P
Food waste compost	1C, 7C, 9C, 10C, 23C, 30C



(a)



(b)

Figure 1. (a) Growth of bacteria producing biofilm (b) Biofilm growth from isolated bacteria

Hydrolysis enzyme screening

Table 2 summarizes the result of hydrolysis enzyme assay for all 13 chosen isolated strains. For cellulase and amylase assay test, after the plate was stained with Gram's iodine, bacteria with positive enzyme assay will have clear hue around the colony on the plate. The result can be seen as shown in Figure 2. In case of lipase enzyme assay test, the reddish – orange colour of the agar will change to yellow in the presence of lipase enzyme as shown in Figure 3. Positive proteolytic enzyme assay test will also show clear hue around the colony on the agar when compared to the milky colour of the agar without the need to stain the agar due to the presence of BCG dye in the agar.




Figure 2. Example for cellulase/amylase enzyme test result after staining




Figure 3. Results of lipase enzyme assay test

Table 2. Summary for the result of hydrolytic enzyme assay test.

Strain	Cellulase assay	Amylase assay	Protease assay	Lipase assay
7	Positive	Positive	Negative	Negative
11	Positive	Positive	Negative	Positive
23	Negative	Negative	Negative	Negative
36	Negative	Positive	Negative	Negative
39	Negative	Positive	Negative	Negative
3P	Negative	Positive	Negative	Negative
19P	Positive	Positive	Negative	Negative
1C	Negative	Negative	Negative	Negative
7C	Negative	Positive	Negative	Negative
9C	Positive	Positive	Negative	Positive
10C	Negative	Positive	Negative	Negative
23C	Positive	Positive	Negative	Positive
30C	Positive	Positive	Negative	Positive

DISCUSSION

All 13 isolated bacteria has been tested for its ability to produce biofilm and it hydrolysis enzyme production. From Figure 1 (a) and (b), there are no conclusive relationship between the growth of bacteria and the amount of biofilm produced by the bacteria. This result is in agreement with previous study [2] that concludes growth rate of bacteria does not correlate with biofilm formation when grown under the same environmental condition. A study done on Salmonella strains also found that different strain have different strength of producing biofilm [15]. The most attractive result from the graph is from strain 1C, which exhibit the highest amount biofilm. Strain 30C have the highest growth but not the highest biofilm producer. The amount of biofilm produced is higher at 48 hr compared to that at 24 hour which are similar to previous studies [2, 15].

All isolated bacteria are negative for protease enzyme assay. This result is good for hydrolysis process, where the absence of protease means that it cannot degrade other enzymes. In a previous research, it was stated that degradation process was slowed down due to the protease breaking down lipase enzyme [14].

Strain number 1C and 23 exhibit negative results for all assay test thus will be excluded for future research. Strain number 11, 9C, 23C, and 30C shows positive results for all cellulase, amylase, and lipase enzyme and will be used for single strain biofilm for simultaneous pre-treatment and enhanced biogas production from POME. The other strains will be mixed for future research.

CONCLUSION

Biofilm producing bacteria are very limited compared to the total bacteria isolated. The biofilm producing bacteria also does not secrete the same enzymes. The depth study need to be done in order to understand the biofilm work for future research on pre-treatment of POME.

ACKNOWLEDGEMENT

This research project was supported by a grant from Fundamental Research Grant (FRG) FRGS15-257-0498. The authors are grateful to the Department of Biotechnology, Kulliyyah of Engineering IIUM for providing the laboratory facilities. The authors would also be very grateful for the supply of food waste compost from IIUM Biolizer team and to West Mill Sdn. Bhd., Sime Darby Research Centre for providing POME and PKC.

REFERENCES

- [1] Langer, S., Schropp, D., Bengelsdorf, F. R., Othman, M., Kazda, M. (2014). Dynamics of biofilm formation during anaerobic digestion of organic waste. *Anaerobe*, 29, 44–51. <http://doi.org/10.1016/j.anaerobe.2013.11.013>
- [2] Sapci, Z. (2013). Bioresource Technology The effect of microwave pretreatment on biogas production from agricultural straws, 128(January 2013), 487–494.
- [3] Taherdanak, M., Zilouei, H. (2014). Improving biogas production from wheat plant using alkaline pretreatment. *Fuel*, 115(October 2016), 714–719. <http://doi.org/10.1016/j.fuel.2013.07.094>
- [4] Tedesco, S., Benyounis, K. Y., Olabi, A. G. (2013). Mechanical pretreatment effects on macroalgae-derived biogas production in co-digestion with sludge in Ireland. *Energy*, 61, 27–33. <http://doi.org/http://dx.doi.org/10.1016/j.energy.2013.01.071>
- [5] Wu, T. Y., Mohammad, A. W., Jahim, J. M., Anuar, N. (2010). Pollution control technologies for the treatment of palm oil mill effluent (POME) through end-of-pipe processes. *Journal of Environmental Management*, 91(7), 1467–1490. <http://doi.org/http://dx.doi.org/10.1016/j.jenvman.2010.02.008>
- [6] Chai, S., Guoa, J., Chaib, Y., Caia, J., Gaoa, L. (2014). Anaerobic treatment of winery wastewater in fixed bed reactors. *Desalination and Water Treatment*, 52(10–12), 1841–1849. <http://doi.org/10.1080/19443994.2013.792008>
- [7] Basri, M. F., Yacob, S., Hassan, M. A., Shirai, Y., Wakisaka, M., Zakaria, M. R., Phang, L. Y. (2010). Improved biogas production from palm oil mill effluent by a scaled-down anaerobic treatment process. *World Journal of Microbiology and Biotechnology*, 26(3), 505–514. <http://doi.org/10.1007/s11274-009-0197-x>
- [8] Szentgyörgyi, E., Nemestóthy, N., Bélafi-Bakó, K. (2010). Anaerobic moving bed biofilm fermenter for biogas production. *Environment Protection Engineering*, 36(4), 117–125.
- [9] Djordjevic, D. (2002). Microtiter plate assay for assessment of *Listeria monocytogenes* biofilm formation. *Applied and Environmental Microbiology*, 68(6), 2950–2958. <http://doi.org/10.1128/AEM.68.6.2950>
- [10] Gohel, H. R., Contractor, C. N., Ghosh, S. K., Braganza, V. J. (2014). A comparative study of various staining techniques for determination of extra cellular cellulase activity on Carboxy Methyl Cellulose (CMC) agar plates. *International Journal of Current Microbiology and Applied Sciences*, 3(5), 261–266.
- [11] Lee, L. P., Karbul, H. M., Citartan, M., Gopinath, S. C. B., Lakshmi Priya, T., Tang, T. H. (2015). Lipase-Secreting *Bacillus* Species in an Oil-Contaminated Habitat: Promising Strains to Alleviate Oil Pollution. *BioMed Research International*, 2015, 1–9. <http://doi.org/10.1155/2015/820575>
- [12] Abdel-Raheem, A., Shearer, C. A. (2002). Extracellular enzyme production by freshwater ascomycetes. *Fungal Diversity*, 11, 1–19.
- [13] Vijayaraghavan, P., Gnana, S., Vincent, P. (2013). A simple method for the detection of protease activity on agar plates using bromocresolgreen dye. *Journal of Biochemical*, 4, 628–630. Retrieved from <http://jbiochemtech.com/index.php/jbt/article/viewArticle/jbt419>
- [14] Bhumibhamon, O., Kopraserstak, A., Funthong, S. (2002). Biotreatment of High Fat and Oil Wastewater by Lipase Producing Microorganisms. *Kasetsart J.(Nat.Sci.)*, 36, 261–267.
- [15] Agarwal, R. K., Singh, S., Bhilegaonkar, K. N., Singh, V. P. (2011). Optimization of microtitre plate assay for the testing of biofilm formation ability in different *Salmonella* serotypes. *International Food Research Journal*, 18(4), 1493–1498.